

Otiderma-Cef[®]

Otic - Dermic Suspension

Antibiotic-antiinflammatory Combination

agrovvetmarket s.a.

FORMULATION

Each 10 mL contains:

Cephalexin monohydrate, (Base).....	200.00 mg
Gentamicin sulphate, (Base).....	100.00 mg
Dexamethasone sodium phosphate.....	0.75 mg
Vitamin A Palmitate.....	10 000 IU
Excipients.....	q.s.ad. 10 mL

INDICATIONS

Otiderma-Cef[®] is indicated for the treatment of otic and dermic problems, associated to allergies and/or susceptible organisms to Gentamicin and/or Cephalexin

Otiderma-Cef[®] is a great support in the treatment of conditions as acute and chronic external otitis, dermatitis and pyoderma. As well as, is indicated as treatment of small cuts, wounds, lacerations, abrasions or when it is necessary to reduce the inflammation and the pain.

The addition of dexamethasone, gives a fast anti-inflammatory effect, reducing also the pain signs and discomfort in the animal, allowing a better arrive of the antibiotics to the infection focus, which gives a major therapeutic effect and effective control.

Besides, **Otiderma-Cef[®]** contains Vitamin A, that acts as epithelial regenerator in animals with infections or inflammation due to pathogens organisms, also reduce the consequences of the infectious process and participate in the tissue integrity.

TARGET SPECIES

The formula is developed for its use in cattle, horses, sheep, goats, pigs, camelids, dogs, cats and rabbits.

SUSCEPTIBLE AGENTS

In vitro, gentamicin acts against a wide variety of gram-positive and gram-negative. Specifically is active against the following organisms: *Alcaligenes* spp., *Citrobacter* spp., *Klebsiella* spp., *Pseudomonas* spp., *Proteus* spp., *Escherichia coli*, *Enterobacter* spp., *Staphylococcus* spp. and *Streptococcus* spp.

The cephalexin is active against: *Staphylococcus* spp. (including penicillin-resistant strains), *Streptococcus* spp., *Corynebacterium* spp., *Pasteurella* spp., *Escherichia coli*, *Proteus* spp., *Micrococcus* spp., *Moraxella* spp., *Actinobacillus ligneresii*, *Actinomyces bovis*, *Haemophilus* spp., *Erysipelothrix rhusiopathiae*, *Clostridium* spp., *Salmonella* spp., *Fusobacterium* spp., *Peptostreptococcus* spp. and *Peptococcus* spp.

ROUTE AND ADMINISTRATION OR APPLICATION WAY

Otic problems: It is recommend, before the use of the product, to clean the auditive canal, dry and verified that there is no rare material or detritus. Stir up the product. The syringe has a large cannula, which offers the possibility to realize a deep application and without causing damage due to its blunt extreme. Apply **Otiderma-Cef[®]** directly in the auditive canal. Once applied, give a slight massage in the base of the ear, in order to the drug adds to the walls and is distribute.

Dermic problems: To use in the skin or mucous membranes, clean the affected area, apply a small quantity of the suspension and distribute uniformly or rub carefully.

DOSAGE

To treat external otitis and other inflammatory conditions of the external ear, fill the external auditive canal with a sufficient quantity of the

suspension until the area to treat is completed.

To its use in skin or mucous membranes, clean the affected area and apply a small quantity of the suspension, distribute or rub smoothly.

Dose intervals: The treatment should be done 1 to 3 times a day, depending of the veterinary doctor criterion.

Minimum duration of the treatment: This varies according the etiology and to the response to the disease curse, it could varies from 2 to 6 weeks, depending on the veterinary doctor criterion.

COLLATERAL EFFECTS AND INCOMPATIBILITIES

It has been not found adverse reactions of any kind in the topical application of the component of **Otiderma-Cef[®]**. However, it should be administered with caution to animals that has shown any allergy form, particularly to penicillin. Those reactions, are very rare, nevertheless if appears, discontinue the treatment and consult to your veterinary doctor.

The gentamicin could produce local reddening. By injectable or oral route, in renal failure and in prolonged treatments, it should be considered as otic and nephrotoxic.

Do not combine with tetracycline, erythromycin, chloramphenicol, due to those antagonize the antibiotic effect.

PRECAUTIONS AND WARNINGS

- Due to the packaging nature and the non/reabsorption of the active ingredient, **Otiderma Cef[®]** present a wide safety margin, making impossible an overdose.
- The product has not show to affect the pregnancy.
- Do not administer the content of the syringe if the protective cap is find broken or damage.
- Preserves in the corresponding dosage unit (syringe) well closed and protected from the direct light exposure.
- Store among 8° to 30° C.
- Keep away from the reach of children.

COMMERCIAL PRESENTATION

Syringe for 15 mL.

Reg. SENASA Perú: F.71.03.N.0014

Otiderma - Cef[®] is a registered trademark of



agrovvetmarket
animal health

Av. Canadá 3792-3798, San Luis. Lima 30 - Perú
Tel.: (511) 435.2323 - 719.0105 Fax: (511) 435.1833
Email: ventas@agrovvetmarket.com - Web: www.agrovvetmarket.com