

FLUNIMINE® | FLUNIX

Injectable Solution

Anti-inflammatory, antipyretic, anti-toxic and non-steroidal analgesic. Specifically indicated for visceral pain





Formulation

Flunixin meglumine 50 mg, excipients q.s. ad 1 mL.

Indications

Cattle and minor ruminants: acute inflammation associated to diseases of the respiratory tract, acute mastitis, post partum paresis, downer cow syndrome, babesiosis and anaplasmosis. Anti-endotoxic in pneumonia, diarrhea, colics and mastitis. Horses: Treatment of choice in horse colic (intestinal anti-inflammatory and anti-endotoxic). Treatment of acute inflammation in cases of musculoskeletal disturbances. Coadjuvant in acute respiratory disease. Pigs: treatment associated to MMA syndrome (mastitis-metritis-agalactia), fever and inflammation of the locomotive system. Dogs and cats: treatment of inflammation and pain associated to illness of respiratory and locomotive systems, and situations of acute pain. In all species: relief of post-surgery and post-traumatic pain. In general, whenever the use of corticoids is contraindicated.

Dosage and Administration

Cattle, sheep, goats, camelids and pigs: 1 - 2 mL for each 45 kg of body weight every 24 hours for 3 - 5 days; horses: 1 mL/45 kg of b.w. every 24 hours for 2 days, the procedure may be repeated in cases of colic after 2 hours or when the symptoms return; dogs: 0.1 mL for each 5 kg of body weight; cats: 0.05 mL each 5 kg of body weight, every 24 hours for 3 - 5 days. Apply through intravenous or intramuscular route.

Commercial Presentation

Bottle x 100 mL,

SENASSA Registry es una referencia informativa y no debe ser considerado como recomendaciones de uso.