Hepaviar® OS

The **liver-boosting** agent for production animals.
Balanced **lipotropic** and **detoxifying** combination.

UNLEASH
THE POTENTIAL

PROTECTS LIVER CELLS FROM TOXIC AGENTS.

PROMOTES THE MOBILIZATION OF FAT ACCUMULATED IN THE LIVER.

HIGHLY ASSIMILABLE INGREDIENTS.

- **+ Silymarin**
- **+ Artichoke Extract (Cynara)**
- **+ N-Acetyl-L-Methionine**
- **+ DL-Methionine**
- **+ L-Carnitine (as Hydrochloride)**
- **+ L-Lysine (as Hydrochloride)**
- **+ Betaine (as Hydrochloride)**
- **+ Choline Chloride**
- **+ Magnesium**
- **+ Sorbitol**



Production Line

Unique formulation

Hepaviar® OS

The **liver-boosting** agent for production animals.

Production Line

Hepaviar® OS is a balanced association of ingredients acting as a lipotropic complex, liver protectant, and detoxifier.

Silymarin promotes protein synthesis, accelerating the liver regeneration process. It adheres to the cell membrane, protecting it from toxic molecules.

Artichoke Extract is a natural liver protectant that facilitates toxin elimination. Its components, cynarin and chlorogenic acid, provide choleretic and cholagogue activity, along with a diuretic effect.

By combining L-carnitine, amino acids, sorbitol, and minerals, the product stimulates correct fat mobilization and proper energy metabolism.



Indications for poultry and swine

- High-energy diets, especially in laying hens.
- Prevention and treatment of Fatty Liver Syndrome.
- *Toxicosis, mycotoxicosis*, and *aflatoxicosis*.
- Acute and chronic hepatitis.
- Preparation for forced molting in poultry.
- Supportive therapy in conditions compromising proper liver function.

Dosage and Administration

Hepaviar® OS must be administered orally mixed with drinking water.

Poultry

Rearing Type	Use	Dosage	Frequency	Treatment Duration
Layers	Preventive	1L / 2000L	Between 28 and 40 weeks of age, every 4 weeks.	3-5 consecutive days
Layers	Preventive	1L / 1000 - 1500L	-	5-7 consecutive days
Broilers	Therapeutic	1L / 2000L	Feed change on the last day of the program.	3-5 consecutive days
Broilers	Therapeutic	1L / 1000 - 1500L	-	5-7 consecutive days

Swine

	Use	Dosage	Frequency	Treatment Duration
Sows Piglets Growers	Preventive	1L / 2000L	Feed change on the last day of the program	3-5 consecutive days
	Therapeutic	1L / 1000 - 1500L	-	5-7 consecutive days

- Cattle and Horses: Direct dose - 1 mL / 5 - 10 kg BW for 5 - 7 days.
- Sheep, Goats, Camelids, Rabbits, Dogs, and Cats: Direct dose – 1 mL / 2.5 – 5 kg BW for 5 – 7 days.

*Consult a specialist before starting any treatment.

Avivet®

For more information:
sales@agrovetmarket.com
+51 (1) 2 300 300 Ext. 101 / +51 922 926 786
www.avivet.net
Av. Canada 3792 - 3798
San Luis, 15021 - Lima - Peru

TÜVRheinland CERTIFIED
Management System
ISO 9001:2015
ISO 27001:2022
www.tuv.com
ID 9105063844