

# Tobrasone®

**NEW**

Combination of Tobramycin with Dexamethasone  
**Sterile Ophthalmic Suspension**

**Effectively treats the main  
infections of the eyeball and  
controls the associated  
inflammation**



# Tobrasone®

## Sterile Ophthalmic Suspension

Tobrasone® is a combination of tobramycin, a bactericidal aminoglycoside antibiotic, and dexamethasone, a potent synthetic glucocorticoid with anti-inflammatory action.

### Composition

Each mL contains:

|                                |      |
|--------------------------------|------|
| Tobramycin Sulfate             | 3 mg |
| Dexamethasone Sodium Phosphate | 1 mg |
| Excipients q.s.p.              | 1 mL |

## Indications for Use:

Treatment and prevention of external eye infections and their adnexa caused by Gram + and Gram - germs, such as *Staphylococcus aureus*, *Moraxella* spp., *streptococci*, and coliforms. Provides an anti-inflammatory effect.

- ✓ Blepharitis, conjunctivitis and/or bacterial keratoconjunctivitis.
- ✓ Corneal infection by *Staphylococcus aureus*.
- ✓ Adjuvant in the treatment of severe intraocular infections.
- ✓ Inflammation of the palpebral, bulbar conjunctiva, cornea, and anterior segment of the eye, when accompanied by bacterial inflammation or when there is a risk of it developing.
- ✓ Corneal abrasions and superficial traumatic lesions of the eyeball.
- ✓ Contaminated keratitis and corneal ulcers caused by susceptible microorganisms, lesions and/or infections in the anterior chamber of the eye.
- ✓ Preventive and post-surgical treatment.



## Administration Routes and Dosage:

Topical ophthalmic route. It is recommended to treat both eyes at the same time due to the risk of a unilateral ocular infection becoming bilateral.

### Moderate Ophthalmic Infections:

Instill 1–2 drops into the affected eye's conjunctival sac every 4–6 hours for the first 24–48 hours. Decrease the frequency as the case evolves, to a frequency of no less than 2 drops every 8 hours. Administer for at least 5 days.

### Severe Ophthalmic Infections:

Instill 2 drops into the affected eye's conjunctival sac every hour until evident improvement is achieved, then continue with 2 drops every 3–4 hours. Administer for at least 7 days.



The doses, frequencies, and duration are for reference and can be adjusted at the veterinarian's discretion.

It is recommended to discontinue therapy at least 48 hours after the infection has been controlled.