

AMINOPLEX® FORTE AC

Injectable Suspension

Amino acids in high concentration



Formulation

Each 100 mL contains:

Dextrose monohydrate.....	5 g	Calcium chloride dihydrate.....	15 mg	Potassium chloride.....	20 mg	Magnesium sulfate trihydrate.....	20 mg	Sodium acetate hydrochloride.....	250 mg	L-Histidine (as hydrochloride).....	27.5 mg	DL-Methionine.....	34 mg	L-Tryptophan.....	34 mg	L-Cysteine (as hydrochloride).....	26.1 mg	L-Threonine.....	68 mg	L-Isoleucine.....	68 mg	L-Arginine (as hydrochloride).....	70.3 mg	L-Phenylalanine.....	102 mg	L-Valine.....	170 mg	L-Lysine (as hydrochloride).....	81.6 mg	L-Leucine.....	136 mg	Monosodium glutamate.....	136 mg	Vitamin B1 (as Thiamine Hydrochloride).....	8.9 mg	Vitamin B2 (as Riboflavin 5-Phosphate).....	2.9 mg	Vitamin B6 (as Pyridoxine Hydrochloride).....	8.2 mg	Vitamin B12 (as Cyanocobalamin).....	5 g	Nicotinamide (Niacinamide).....	150 mg	Excipients.....	q.s.	to.....	100 mL
---------------------------	-----	---------------------------------	-------	-------------------------	-------	-----------------------------------	-------	-----------------------------------	--------	-------------------------------------	---------	--------------------	-------	-------------------	-------	------------------------------------	---------	------------------	-------	-------------------	-------	------------------------------------	---------	----------------------	--------	---------------	--------	----------------------------------	---------	----------------	--------	---------------------------	--------	---	--------	---	--------	---	--------	--------------------------------------	-----	---------------------------------	--------	-----------------	------	---------	--------

Indications

As an immediate source of amino acids, dextrose, B-complex vitamins, and electrolytes for the treatment of acute convalescent states in animals that are debilitated or subjected to intense exertion, stress, shock, dehydration, chronic diseases, or parasitism; and as supportive therapy in cases of severe dehydration caused by vomiting or diarrhea, where it aids in replenishing lost electrolytes and proteins (in such cases, oral or parenteral fluid therapy is recommended).