



Soluble Powder

Synergistic Antibiotic Combination of Wide Spectrum and Long Action





Formulation

Combiótico ® L.A. SPI 3.6 g Bottle 1: Penicillin G Benzathine 2 000 000 I.U., dihydrostreptomycin sulfate equivalent to 2 g base. Bottle 2: Sterile diluent 8 mL. Each reconstituted mL contains: Benzathine Penicillin G 200,000 IU, dihydrostreptomycin sulfate equivalent to 0.2 g base, excipients q.s.p.1 mL. Combiótico ® L.A. SPI 13.5 g Bottle 1: Penicillin G Benzathine 7,500,000 I.U., dihydrostreptomycin sulfate equivalent to 7.5 g base. Bottle 2: Sterile diluent 30 mL. Each reconstituted mL contains: Benzathine Penicillin G 200,000 IU, dihydrostreptomycin sulfate equivalent to 7.5 g base.

Indications

Treatment and prevention of local or systemic infections caused by germs susceptible to penicillin and / or dihydrostreptomycin, provided that a long-acting effect is required (spaced doses)

Dosage and Administration

Intramuscular injectable route (in intramuscular and subcutaneous canines), repeat after 72 hours if necessary.Cattle, sheep, goats, horses, South American camelids and pigs: the recommended dose is 1 mL for every 10-20 kg of live weight. In young animals: Calves, lambs, foals, piglets, the dose is 1 - 2 mL / 25 kg. Canines and felines: the recommended dose is 1 - 2 mL / 25 kg of live weight.

Commercial Presentation

Bottle of 3.6g, Bottle of 8mL of diluent, 30mL bottle of diluent, 13.5mL bottle of diluent,

SENASA Registry

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.