COXURIL T® 50 OS



Oral Solution

Coccidicide Effective Against All Eimeria Species Including Those Resistant to Other Compounds - High Concentration

Formulation

Each 100 mL contains: Toltrazuril 5 g, excipients q.s.p. 100 mL

Indications

Treatment, prevention, and control of infections caused by coccidia. - Broilers, pullets, and young hens: Eimeria acervulina, E. brunetti, E. maxima, E. mitis, E. necatrix, E. embati, E. mivati, and E. tenella. - Turkeys: E. adenoides and E. meleagrimitis. - Geese: E. anseris and E. truncata. - Pigeons: E. columbarum and E. labbeana. Its broad spectrum includes all coccidia species of economic importance in poultry.

Dosage and Administration

Orally, with drinking water in birds for two consecutive days. In general: 7 mg of toltrazuril/kg of body weight, which is equivalent to 1 L of the product/2,000 liters (1 mL/2 L) drinking water (25 ppm)* * Other specific doses. Chickens: 10 to 15 ppm against: Eimeria acervulina, E. brunetti, E. maxima, E. tenella, E. mivati and E.necatrix. Geese: 20 ppm against: E. anseris and E. truncata. Pigeons: 5 mg/kg against: E. columbarum and E. labbeans. Turkeys: 15 to 25 ppm against: E. adenoids and E. meleagrimitis.

Commercial Presentation

Bottle x 100 mL, Bottle x 250 mL, Bottle x 1 L,

SENASA Registry

F.08.32.I.1121

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.

Copyright © 2026 Agrovet Market S.A.