DIFLOVET® 10

Injectable Solution

Second-generation Fluoroquinolone antibacterial





Formulation

Difloxacin hydrochloride 100 mg, excipients s.q.ad. 1 mL

Indications

For the treatment and prevention of infections caused by difloxacin sensitive microoganisms, basically of the respiratory, genito-urinary and digestive tract, skin and ear canal, bovine respiratory disease, colibacillosis, salmonellosis, pneuomnia, bronchopneumonia, mycoplasmosis, dermatitis, mastitis-metritis-agalactia, rhinitis, among others.

Dosage and Administration

Cattle, sheep, camelids, goats and swines: 1 mL/40 kg of b.w. for 3 to 5 days; dogs: 1 mL/20 kg in a single subcutaneous injection, followed by oral treatment of difloxacin in tablets (5 mg/kg of b.w.) for a minimum of 5 days; poultry: 0.1 mL/kg for 3 to 5 days. Apply through subcutaneous route; the intramuscular route may be used only in swine and poultry.

Commercial Presentation

Bottle x 100 mL,

SENASA Registry

SENASA Perú: F.03.01.N.1031

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.

Copyright © 2026 Agrovet Market S.A.