

DUOCORTYL® AC

Injectable Suspension

Anti-inflammatory - Immunosuppressant -
Antiallergic - Antitoxic - Gluconeogenetic



Formulation

Each 1 mL contains: Prednisolone (as acetate) 7.5 mg, Dexamethasone (as sodium phosphate) 2.5 mg, Excipients q.s.ad 1 mL.

Indications

- Non-infectious inflammatory processes, especially musculoskeletal inflammations (arthritis, osteoarthritis, tendovaginitis, bursitis, dislocations, myositis and sprains among other injuries). - Acute infectious diseases -associated with antibiotic therapy-: osteoarthritis, lymphangitis, arthritis, bronchopneumonia, septicemia, cholinerotoxemia, mastitis and infections of the urogenital tract, otitis, ocular disorders¹, dermatitis, phlegmon among others. - Immunosuppression in cases of allergic conditions such as anaphylaxis, asthma, non-specific skin conditions (atopic dermatitis, urticaria, pruritus), laminitis, stings, snake bites, drug allergies and in cases of autoimmune diseases: lupus erythematosus, rheumatoid arthritis and others; as well as in tumor processes and other conditions. - Emergency therapies, such as shock and circulatory collapse (eg anaphylaxis, poisoning, etc.) - Conditions of physical stress (eg febrile disorders, trauma, surgery, weakness, burns) - Adrenal endocrine disorders, in patients with -known adrenal insufficiency or when there are doubts about their adrenocortical reserves.

Dosage and Administration

Intramuscular or subcutaneous injection route. Shake before using. Canines: 0.2 - 0.6 mL, felines: 0.1 - 0.3 mL. Doses can be adjusted according to the pathology to be treated, according to the veterinarian's criteria. The desired result is usually achieved after just one application. If maintenance treatment is necessary after the initial therapeutic effect has been achieved, it will be carried out at reduced doses, set at the discretion of the veterinary medical professional. Do not administer for more than 5 days.



Commercial Presentation Este es un producto de referencia informativa y no debe ser considerado como recomendaciones de uso.