ECTOMETHRIN® 200

Emulsifiable Concentrate

Pyrethroid ectoparasiticide





Formulation

Cypermethrin 200 mg; excipients q.s. ad 1 mL.

Indications

For the treatment and control of ticks, fleas, mites, lice and flies, among others. Its residual action lasts from 3 to 4 weeks. It can be used in immersion baths, sprinkling or spraying of buildings and facilities.

Dosage and Administration

For immersion baths: Foot bath: 1 L/1250 L of fresh or salty water. Replenishing and restrengthening the solution: 1.5 L/1250 L of water. Both operations to be performed when the level in the bath is reduced in 10% of its original level. Dry restrengthening: 0.5 L/1250 L of water consumed in the bath. This is done in order to not having to replenish the water consumed. For sprinkling animals: 1 L/1250 L of water (apply the final sprinkling emulsion using approximately 1 L/animal). Dogs: 1 mL/1.25 L of water. For spraying in buildings and facilities: 1 mL/1.5 to 2.5 L of water (1 L of the final emulsion per 10 to 15 m2 of surface to be treated).

Commercial Presentation

Bottle x 20 mL, Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Bottle x 1 L,

SENASA Registry

SENASA Perú: F.07.08.N.0320

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.

Copyright © 2026 Agrovet Market S.A.