

HISTACORT® AC

Injectable Suspension

Potent Synergistic Anti-allergic Dual Mechanism of Action Association



Formulation

Each mL contains: Prednisolone (as acetate)..... 10
mg Chlorpheniramine maleate..... 4 mg
Excipients..... q.s.ad..... 1 mL

Indications

Symptomatic treatment of pathologies associated with histamine release and as a corticosteroid anti-inflammatory whenever an antihistamine effect is required. Allergic reactions in general, anaphylaxis¹, allergic dermatosis/dermatitis, atopic dermatitis, contact dermatitis, stinging dermatitis, pruritus² (regardless of its cause) and allergic inflammation associated with allergic, parasitic, pustular and non-specific dermatitis and otitis. Drug and food reactions, itching, urticaria. Respiratory allergies, allergic rhinitis, pollinosis, ocular allergies, inflammatory ophthalmic disorders³ when systemic corticotherapy is indicated, allergic bronchitis (asthma), shock states. Tracheobronchitis, bronchitis, including all allergic bronchitis. Bronchospasm, coadjuvant in the treatment of arthritic and rheumatic diseases, arthropathies, autoimmune diseases, bursitis among others; whenever a corticoid effect associated to the antihistamine is required.

Dosage and Administration

By intramuscular or subcutaneous injection. Dogs and cats, at the rate of: prednisolone: 1 mg/kg and chlorpheniramine: 0.4 mg/kg of body weight. In practice: 1 mL/10 kg body weight. Doses can be adjusted according to the veterinarian's criteria. Histacort® AC can be administered to animals suffering from acute or chronic bacterial infections, provided that the infection is controlled with appropriate antibiotic or chemotherapeutic agents.

Commercial Presentation



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.