IMIDOX® 120

Injectable Solution

Antiprotozoal hematicide Agent





Formulation

Imidocarb dipropionate 120 mg, excipients .q.s.ad. 1 mL

Indications

For the treatment and prevention (before the introduction of livestock to areas of risk) against piroplasmosis, anaplasmosis (both also known as "tick fever"), ehrlichiosis, hepatozoonosis and cytauxzoonosis.

Dosage and Administration

Piroplasmosis: cattle, sheep and goats: 1 ml/100 kg; horses: 2 mL/100 kg; dogs and cats: 0.25 mL/10 kg. Anaplasmosis and piroanaplasmosis: cattle, sheep and goats: 2.5 mL/100 kg; Ehrlichiosis: dogs: 0.5 mL/10 kg of b.w.; Cytauxzoonosis: cats: 0.2 mL/10 kg of b.w. Apply through intramuscular route in horses, and preferably subcutaneously in the dorsal area of cattle, sheep, goats, dogs and cats.

Commercial Presentation

Bottle x 20 mL, Bottle x 100 mL,

SENASA Registry

SENASA Perú:F.08.01.N.0478.

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.

Copyright © 2026 Agrovet Market S.A.