

MITRAZYDE® 200

Emulsifiable Concentrate

High-Concentration Acaricide and Tickicide



Formulation

Each 100 mL contains: Amitraz..... 20 g Excipients q.s.
to.....100 mL

Indications

Mitrazyde® 200 is indicated for the prevention and treatment of infestations by ticks, mange mites (sarcoptic, demodectic, and notoedric), lice, keds, among other ectoparasites.

Dosage and Administration

Mitrazyde® 200 is administered topically (externally) via immersion baths or spraying, and by topical application to localized lesions. Dilute according to the dosage and shake until a homogeneous white mixture is obtained. Immersion Baths Cattle, South American camelids, sheep, and goats. Initial dip: Dilute the product at a ratio of 1–1.25 L per 1000 L of water (200–250 ppm or mg/L). Reinforcement and top-up: 2.5 L per 1000 L of water (500 ppm). Apply when the water level in the pond drops below 1000 L. Spray dip (spray sleeve, backpack sprayer, or hand sprayer): - Cattle: Add 25 mL of the product per 20 L of water (250 ppm), mix well, and thoroughly bathe the cattle. Demodectic or sarcoptic mange: 50 mL per 20 L of water (500 ppm). Use 4–5 L of the mixture to bathe 1 adult bovine or 2 calves. - Swine: Add 40 mL of the product per 20 L (400 ppm) of water

Commercial Presentation

Bottle x 30 mL, Bottle x 100 mL, Bottle x 250 mL, Bottle x 500 mL, Bottle x 1 L,

SENASA Registry

Reg. SENASA Peru: F.087.046.I.00646



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.