

NUTROMIX® LYSINE+

Oral Gel

High Availability L-Lysine Nutritional Supplement



Formulation

Each 100 g contain:L-Lysine HCI	8.58
gExcipients c.s.p 100g	

Indications

Conditions that respond to lysine, such as upper respiratory problems or Feline Herpes Virus (FHV-1). In the acute phase of FHV-1, to reduce the severity of symptoms, accelerate healing and minimize the ability to of the sick cat to spread contagion. As FHV-1 infection is chronic, L-lysine is especially indicated as preventive maintenance in order to improve the quality of life of cats with chronic symptoms, avoid relapses and minimize the risk of contagion to other cats in the home.

Dosage and Administration

Administer orally. Acute phase: Cats: 500-1,000 mg of L-Lysine HCI / animal (6-12 g, 10-20 cm or 1-2 teaspoons). Kittens under 6 months: half the dose. Once or twice a day. Chronic or latent phase: Adult cats, 500 mg (6g, 10 cm or 1 teaspoon); Kittens: half the dose. Once a day. As a preventive: daily or at least during periods of stress (eg, entry of a new cat into the house, pregnancy, change of season, etc).

Commercial Presentation

SENASA Registry

Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.