

PLEUROTIL® 250 OS

Oral Solution

Macrolide Antibacterial with Selective Pulmonary Affinity



Formulation

Each 100 mL contain: Tilmicosin 25 g, excipients q.s. ad 100 mL.

Indications

Chicken and turkeys: respiratory infections caused by *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Ornithobacterium rhinotracheale*, *Pasteurella multocida* and other microorganisms sensitive to tilmicosin. Pigs: treatment of respiratory infections caused by *Pasteurella multocida*, *Pasteurella haemolytica*, *Mycoplasma* spp., *Bordetella bronchiseptica*, *Staphylococcus aureus*, *Erysipelothrix rhusiopathiae*, *Klebsiella* spp., *Haemophilus parasuis*, *Salmonella* spp., *Pseudomonas* spp., and *Streptococcus* spp., among other microorganisms sensitive to tilmicosin. Calves: cattle respiratory syndrome associated to *Mannheimia haemolytica* and *Pasteurella multocida*. Treatment of interdigital necrobacillosis.

Dosage and Administration

By oral route, with drinking water. Poultry: 300 mL/1000 L, during 3 days. Pigs: 800 mL/1000 L during 5 days, in drinking water. Cattle and minor ruminants: 1 mL/20 kg of body weight during 3 - 5 days.

Commercial Presentation

Bottle x 100 mL, Bottle x 1 L,

SENASA Registry

SENASA Perú: F.03.32.I.1816



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.