

SURALAN® QUATTRO® SMALL

Soft Chews

Multiple Action Ecto-Endoparasitic, Low Dose, 5 Weeks of Protection



Formulation

Each 2000 mg chewable tablet contains: Sarolaner (S-Sarolaner): 20.00 mg Moxidectin: 0.24 mg Pyrantel (as embonate): 50.00 mg Arpraziquantel (R-Praziquantel): 25.00 mg Excipients and flavorings q.s. to 1 tablet

Indications

Suralan® Quattro® is indicated for the treatment of dogs with, or at risk of, mixed parasitic infestations caused by endoparasites and ectoparasites, provided that simultaneous action is required against fleas, ticks, mites, nematodes (including gastrointestinal, pulmonary, ocular and heart) and cestodes (tapeworms), also preventing dirofilariosis and angiostrongylosis.

Dosage and Administration

Route of Administration: Oral Dog Dosage: Dog Weight: 5 - 10 kg Dosage: 1 tablet every 35 days (5 weeks) Dosage per active ingredient: Sarolaner: 2-4 mg/kg Moxidectin: 0.024-0.048 mg/kg Pyrantel: 5-10 mg/kg Arpraziquantel: 2.5-5 mg/kg Administration Method: Can be administered with or without food (before or after feeding). Palatable, soft, and chewable tablet for easy administration. If the dog does not consume it voluntarily, it can be administered with food or by placing the tablet deep at the back of the tongue. Do not divide the tablets. Frequency according to condition: Fleas and ticks: once every 35 days during the season. Sarcoptic mange: monthly administration for at least 2 consecutive months. Demodectic mange: once a month during At least 3 consecutive months (continue until negative skin scrapings are obtained on at least 2 consecutive occasions with a 1-month interval) Otodectic mange: single dose sufficient (veterinary examination recommended; veterinary check-up at 28 days) Gastrointestinal nematodes: single dose for active infestations, then monthly as a preventative Dirofilariosis and angiostrongylosis: monthly intervals in endemic areas (continue until at least 1 month after the last mosquito exposure) Angiostrongylosis (treatment): single dose (veterinary check-up at 30 days) Cestodes: apply once as a single treatment, then monthly as a preventative Special considerations: Can be used from 8 weeks of age and 1.25 kg of weight Puppies should be weighed regularly for dose adjustment Do not administer at intervals of less than 30 days Treat all animals in the household at the same time for flea control