TYLAMOX® WS

Soluble Powder

Broad-spectrum Macrolide Beta-lactam
Synergic-additive Antibiotic Association





Formulation

Each 100 g contains: Tylosin as tartrate 50 g, Amoxicillin as trihydrate 10 g, excipients q.s. ad 100 g.

Indications

Treatment and prevention of infections caused by sensitive germs: infections of the respiratory, digestive and urogenital systems, infections of the skin and soft tissues, and prevention of secondary bacterial infections. Poultry: CRD and other respiratory diseases, salmonellosis, colibacillosis, cholera, necrotic enteritis, staphylococcosis and streptococcosis, coryza, omphalitis and infection of the vitelline sac, sacculitis, sinusitis, synovitis, dermatitis. Pigs: enzootic pneumonia and other respiratory diseases, dysentery, atrophic rhinitis, edema disease, colibacillosis, salmonellosis, diarrheas, corinebacteriosis, erysipelosis, leptospirosis, dermatitis and others. Pre ruminants: respiratory and enteric infections, diarrheas, enterotoxemia.

Dosage and Administration

During 3 to 5 days: Poultry: Preventive: 1 kg/2000 L of water (or in 1000 kg of feed). Preventive for mycoplasmosis and therapeutic: 1 kg/1000L of water (or in 500 kg of feed). Pigs: Preventive: 1 kg/4000 L of water (or in 2000 kg of feed). Therapeutic: 1 kg/2000 - 4000 L of water (or in 1000 - 2000 kg of feed). Pre-ruminants: Therapeutic: 2 times a day at a rate of 1 g/20 kg each time (in water, milk or feed). Preventive:half the dose.

Commercial Presentation

Sachet x 100 g, Sachet x 500 g, Sachet x 1 kg, Sachet x 15 g,

SENASA Registry

~~Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.~~

SENASA Perú: F.82.70.N.0023

Copyright © 2026 Agrovet Market S.A.