

TYLODOX® WS

Soluble Powder

Broad-spectrum Bacteriostatic - Additive Antibiotic Combination



Formulation

Each 100 g contains: Tylosin as tartrate 10 g, Doxycycline as hyclate 10 g, excipients q.s. ad 100 g.

Indications

Treatment and prevention of infections caused by germs sensitive to tylosin and/or doxycycline. Among them: infections of the respiratory, digestive and urogenital systems, such as: mycoplasmosis, CRD, coryza, colibacillosis, omphalitis and infection of the vitelline sac, avian cholera, enteritis and diarrhea, necrotic enteritis, sacculitis, sinusitis, synovitis, staphylococcosis, streptococcosis, corinebacteriosis, hexa-mitiasis and leptospirosis. Prevention of secondary bacterial infections following viral infections, stress conditions, vaccinations, re-lodging, beak trimming, transportation, feed and climate changes.

Dosage and Administration

Poultry and pigs: Preventive: 1 kg/2000 L of water (or in 1000 kg of feed) during 3 - 5 days. Treatment: 1 kg/1000 - 1500 L of water (or in 500 - 750 kg of feed) during 3 - 5 days. Pre-ruminants: Therapeutic: 2 times a day at a rate of 1 g/20 kg each time during 3 - 5 days (in water, milk or feed). Preventive: half the dose.

Commercial Presentation

Sachet x 100 g, Sachet x 500 g, Sachet x 1 kg, Sachet x 15 g,

SENASA Registry

SENASA Perú: F.03.70.I.1695



Este contenido es una referencia informativa y no debe ser considerado como recomendaciones de uso.